arXiv:gr-qc/0501100v1 31 Jan 2005

Geometric description of lightlike foliations by an observer in general relativity

V. J. Bolós

Dpto. Matemáticas, Facultad de Ciencias, Universidad de Extremadura.
Avda. de Elvas s/n. 06071–Badajoz, Spain.
e-mail: `vjbolos@unex.es`

January 31, 2005

Abstract

We introduce new concepts and properties of lightlike distributions and foliations (of dimension and co-dimension 1) in a space-time manifold of dimension n, from a purely geometric point of view. Given an observer and a lightlike distribution Ω of dimension or co-dimension 1, its lightlike direction is broken down into two vector fields: a timelike vector field U representing the observer and a spacelike vector field S representing the relative direction of propagation of Ω for this observer. A new distribution Ω_U^- is defined, with the opposite relative direction of propagation for the observer U. If both distributions Ω and Ω_U^- are integrable, the pair $\left\{\Omega, \Omega_U^-\right\}$ represents the wave fronts of a stationary wave for the observer U. However, we show in an example that the integrability of Ω does not imply the integrability of Ω_U^-.

1 Introduction

It is known that foliations theory is an appropriate framework to study wave fronts [5]. For example, in symplectic mechanics, the evolution of a dynamical system is studied by the leaves of a foliation associated to a symplectic manifold $(\mathcal{U}, \sigma)$: this dynamical system has a space of evolution with the structure of a presymplectic manifold $(\mathcal{V}, \sigma_{\mathcal{V}})$, whose characteristic foliation, $ker\sigma_{\mathcal{V}_\flat}$, describes [5] the evolution of $(\mathcal{U}, \sigma)$. In a particular case, in Special Relativity, for any observer the evolution of a free massless elementary particle is described by 2-planes moving in the normal direction at the speed of light [5] (i.e. lightlike moving wave fronts). This result that holds in the Minkowski space-time is not trivially generalized to General Relativity [3].

In this paper we are only interested in the dynamics of the wave fronts as leaves of a foliation. So, we introduce new concepts and properties of lightlike distributions and foliations (of dimension and co-dimension 1) in order to improve this description of the wave fronts, from a purely geometric point of view.

In Sec. 2, 3, 4 we work with distributions to make the study as general as possible, with the aim of making the results applicable to a wider range of situations. In Sec. 2, given an observer and a lightlike distribution Ω, its lightlike direction is broken down into two vector fields: a timelike vector field U representing the observer and a spacelike vector field S representing the relative direction of propagation of Ω for this observer. In Sec. 3 we study the change of observer, and in Sec. 4 we define a new distribution Ω_U^- with the opposite relative direction of propagation for the observer U. In Sec. 5 we give more properties about the change of observer, and in Sec. 6 we study the case that both distributions Ω and Ω_U^- are integrable. However, we show in Example 9 that the integrability of Ω does not imply the integrability of Ω_U^-. In Sec. 7 we give some physical interpretations and applications of some obtained results. For example, if Ω and Ω_U^- are integrable, the pair $\left\{\Omega, \Omega_U^-\right\}$ represents a stationary wave for the observer U. Finally, we apply some results to give an interpretation of the light aberration in General Relativity.

We will work on an n-dimensional space-time manifold M $(n > 2)$ with metric g given by $ds^2 = g_{ij}dx^i dx^j$ and $c = 1$. A list of vector fields inside *span* () will denote the subbundle generated by these vector fields (called *distribution*). Usually, a distribution of dimension p is called a p-distribution. All adapted bases of distributions are local. A distribution that has an integral submanifold (leaf) in every point is a *foliation.* We will say that a future pointing timelike unitary vector field is an observer (identifying it with the 4-velocities of a congruence of timelike world lines), and usually we will denote it by U. All the causal vectors (lightlike or timelike) will be considered future-pointing. We will denote by $\Omega^\perp$ the orthogonal distribution of Ω. If v is a vector, $v^\perp$ denotes the orthogonal subspace of v. Moreover, if v is a spacelike vector, $\|v\|$ denotes the module of v.

2 Introducing the concept of "basis associated to an observer"

A lightlike 1-distribution is a field of 1-dimensional lightlike subspaces and a lightlike $(n-1)$-distribution is a $(n-1)$-dimensional distribution of subspaces whose normal is a lightlike vector field, which must therefore be contained in the distribution itself. So, both 1-distributions and $(n-1)$-distributions are completely determined by a lightlike vector field N representing the "lightlike direction" of the distribution (up to scale factors). Moreover, two distributions Λ and Ω (of dimension and co-dimension 1, respectively) are orthogonal if and only if they have the same lightlike direction. In this case, the 1-distribution is contained in the $(n-1)$-distribution: $\Lambda \perp \Omega \Longleftrightarrow N \in \Lambda \subset \Omega$.

In each case (1-distribution and $(n-1)$-distribution) this lightlike direction can always be expressed as the sum of an observer U and an orthogonal unitary spacelike vector field S which represents the relative direction of N for this observer. So, a basis of a lightlike 1-distribution Λ is given by

$$\{S+U\} \tag{1}$$

where $S \in U^{\perp}$ and $\|S\| = 1$. We will say that (1) is the *U-basis of* Λ (also known as *basis of* Λ *associated to* U), since it is completely determined by U. Moreover, we will say that S is the *direction of the relative velocity of* Λ *observed by* U.

Example 1 *In the Minkowski space-time (with $n = 4$), expressing the metric g in rectangular coordinates, we consider the 1-distribution Λ generated by*

$$\left\{\frac{\partial}{\partial z} + \frac{\partial}{\partial t}\right\}. \tag{2}$$

If we consider the observer $U = \frac{\partial}{\partial t}$, then (2) is the U-basis of Λ of the form $\{S+U\}$ where $S = \frac{\partial}{\partial z}$ is a unitary spacelike vector field orthogonal to U. But if we consider the observer $U' = \sqrt{2}\left(\frac{\partial}{\partial t} + \frac{1}{\sqrt{2}}\frac{\partial}{\partial x}\right)$, then (2) is not the U'-basis of Λ. It is given by $\{S'+U'\}$ where $S' = \sqrt{2}\left(\frac{\partial}{\partial z} - \frac{1}{\sqrt{2}}\frac{\partial}{\partial x}\right)$ is a unitary spacelike vector field orthogonal to U'. Logically, $S+U$ and $S'+U'$ represents the same lightlike direction (i.e. they are proportional).

On the other hand, a lightlike $(n-1)$-distribution Ω can therefore only contain $(n-2)$ independent spacelike vector fields since timelike vectors cannot be orthogonal to lightlike vectors by simple special relativistic considerations. Given an observer U, a basis of Ω can therefore be chosen to be $S+U$ (a representative of the lightlike direction of Ω) and $(n-2)$ independent spacelike unitary vector fields, $X_1, \ldots, X_{n-2}$:

$$\{X_1, \ldots, X_{n-2}, S+U\} \tag{3}$$

such that $X_1, \ldots, X_{n-2}, S$ are an orthonormal basis of $U^{\perp}$. So $X_1, \ldots, X_{n-2}$ form an orthonormal basis of $\Omega \cap U^{\perp}$ and $\{X_1, \ldots, X_{n-2}, S, U\}$ form an orthonormal basis of TM (called tetrad in the case $n = 4$, see [2]). We will say that (3) is a *U-basis of* Ω (also known as a *basis of* Ω *associated to* U). Moreover, we will say that S is the *direction of the relative velocity of* Ω *observed by* U. $X_1, \ldots, X_{n-2}$ depends on U but they are not completely determined (we can make infinite choices, since they form an orthonormal basis of $\Omega \cap U^{\perp}$, i.e. they complete with S an orthonormal basis of $U^{\perp}$).

Example 2 *In the Minkowski space-time, expressing the metric g in rectangular coordinates, we consider the 3-foliation Ω generated by*

$$\left\{\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} + \frac{\partial}{\partial t}\right\}. \tag{4}$$

If we consider the observer $U = \frac{\partial}{\partial t}$*, then (4) is a* U*-basis of* Ω *of the form* $\{X_1, X_2, S+U\}$ *where* $X_1 = \frac{\partial}{\partial x}$, $X_2 = \frac{\partial}{\partial y}$, $S = \frac{\partial}{\partial z}$ *are an orthonormal basis of* $U^{\perp}$*. But if we consider the observer* $U' = \sqrt{2}\left(\frac{\partial}{\partial t} + \frac{1}{\sqrt{2}}\frac{\partial}{\partial x}\right)$*, then (4) is not a* U'*-basis of* Ω*. It is given by* $\{X_1', X_2', S'+U'\}$ *where* $X_1' = \frac{\partial}{\partial x} + \frac{1}{\sqrt{2}}\left(\frac{\partial}{\partial z} + \frac{\partial}{\partial t}\right)$, $X_2' = \frac{\partial}{\partial y}$, $S' = \sqrt{2}\left(\frac{\partial}{\partial z} - \frac{1}{\sqrt{2}}\frac{\partial}{\partial x}\right)$ *are an orthonormal basis of* $(U')^{\perp}$*. It is important to remark that* S' *is completely determined by* U'*, but we can choose* X_1'*,* X_2' *to complete this orthonormal basis. Logically,* $S+U$ *and* $S'+U'$ *represent the same lightlike direction (i.e. they are proportional), and the subspace* $\operatorname{span}(X_1, X_2)$ *is not the same as the subspace* $\operatorname{span}(X_1', X_2')$*.*

So, the vector fields of a basis associated to an observer describe the distribution from the point of view of this observer.

3 Change of observer

With this notation, given a lightlike distribution (of dimension or co-dimension 1) and two different observers U, U', a change of observer is a change from a U-basis to a U'-basis. Now we are going to study this change: U' can be always written in the form

$$U' = \gamma (U + vX) \tag{5}$$

where $X \in U^{\perp}$, $\|X\| = 1$, v is a differentiable function such that $0 < v < 1$ and $\gamma = \frac{1}{\sqrt{1-v^2}}$. It is easy to prove that this decomposition is unique. We will say that v is the *module of the relative velocity of* U' *observed by* U and X is the *direction of the relative velocity of* U' *observed by* U [1]. So, γ is the *gamma factor* corresponding to the velocity v.

Let $\{S+U\}$ and $\{S'+U'\}$ be the U-basis and the U'-basis, respectively, of the same lightlike 1-distribution Λ. Then, it is easy to prove that

$$S' = \frac{1}{\gamma\left(1 - vg\left(X, S\right)\right)}\left(S+U\right) - \gamma\left(U + vX\right). \tag{6}$$

The fact that S' is different from S is the *aberration* effect.

In the same way, let $\{X_1, \ldots, X_{n-2}, S+U\}$ and $\{X_1', \ldots, X_{n-2}', S'+U'\}$ be a U-basis and a U'-basis, respectively, of the same lightlike $(n-1)$-distribution Ω. Then, S' is given by (6) too and, by direct calculus (orthonormalization of $\{X_1, \ldots, X_{n-2}, S', U'\}$), we obtain that a choice of X_i' can be

$$X_i' = X_i + \frac{vg\left(X, X_i\right)}{1 - vg(X, S)}\left(S+U\right)$$

for $i = 1, \ldots, n-2$.

4 Equivalence relations between distributions and observers

In this Section, we are going to define equivalence relations between distributions (see Definition 3) and between observers (see Definition 4).

Definition 3 *Given an observer U, let Ω and Ω' be two lightlike p-distributions of the same dimension ($p = 1, n-1$) with lightlike directions $S+U$ and $S'+U$, respectively. We will say that these p-distributions are equal up to orientations for the observer U, if and only if $S' = \pm S$. It will be denoted by $\Omega \overset{U}{=} \Omega'$ u.t.o.*

In this case, the relative directions of the lightlike directions of these p-distributions are the same or the opposite for this observer.

Given Ω a lightlike p-distribution ($p = 1, n-1$) and U an observer, we can build another lightlike p-distribution with the opposite relative direction of propagation for this observer, and it will be denoted by Ω^-_U. So, if $p = 1$ and $\{S+U\}$ is the U-basis of Ω, then $\{-S+U\}$ is the U-basis of Ω^-_U. On the other hand, if $p = n-1$ and $\{X_1, \ldots, X_{n-2}, S+U\}$ is a U-basis of Ω, then $\{X_1, \ldots, X_{n-2}, -S+U\}$ is a U-basis of Ω^-_U. In both cases ($p = 1, n-1$) we have that Ω^-_U is well defined and obviously $\left(\Omega^-_U\right)^-_U = \Omega^{--}_{UU} = \Omega$.

It is not difficult to prove that $\Omega \overset{U}{=} \Omega'$ u.t.o. is equivalent to $\Omega^\perp \overset{U}{=} \Omega'^\perp$ u.t.o., and it is clear that

$$\left(\Omega^-_U\right)^\perp = \left(\Omega^\perp\right)^-_U . \tag{7}$$

Summing up, given an observer U, the relation "be equal up to orientations for the observer U" defines a quotient space on the set of the lightlike distributions of dimension or co-dimension 1. This quotient space is the framework to study stationary waves, as we will see in Sec. 7.2. Each equivalence class is defined by two distributions, Ω and Ω^-_U, i.e.:

$$\Omega \overset{U}{=} \Omega' \quad u.t.o. \quad \Longleftrightarrow \quad \Omega' = \Omega \quad or \quad \Omega' = \Omega^-_U. \tag{8}$$

From a geometrical point of view, the lightlike directions of Ω and Ω^-_U are in a 2-plane which contains the observer U (see Fig. 1 and Fig. 2).

Let us now introduce the concept of "observers Ω-related":

Definition 4 *Given a lightlike distribution Ω (of dimension or co-dimension 1) and two observers, U and U', we will say that these observers are Ω-related if and only if $\Omega^-_U = \Omega^-_{U'}$. It will be denoted by $U \overset{\Omega}{\approx} U'$.*

Then, two observers are Ω-related if and only if they are coplanar with the lightlike direction of Ω (see Fig. 3). So, if these observers are observing Ω, there is not any aberration effect between them, as we will see in Sec. 7.1.

Properties of observers Ω-related are compiled in the next theorem:

Theorem 5 *Given a lightlike p-distribution Ω ($p = 1, n-1$) and two observers, U and U', the following properties are equivalent (see Fig. 3 and Fig. 4):*



Figure 1: The lightlike directions of Ω and Ω_U^- are represented as N and N', respectively (lightlike vectors in the future lightcone), and U is represented as a vector inside the lightcone, since it is future-pointing timelike unitary



Figure 2: Figure 1 could be represented schematically in a section of the lightcone: lightlike directions are represented as points of the circle (that represents the section), observers are represented as points inside the circle and the 2-plane defined by the lightlike directions of Ω and Ω_U^- is represented as a straight line joining both lightlike directions. Since a lightlike direction determines the lightlike p-distribution ($p = 1, n-1$), the lightlike direction is represented by the name of the distribution which determines. So, the lightlike direction of Ω is represented as "Ω" and the lightlike direction of Ω_U^- is represented as "Ω_U^-"


$\Omega=\Omega^{--}_{UU'}=\Omega^{--}_{U'U}$
$=\Omega^{--}_{UU}=\Omega^{--}_{U'U'}$
U
U'
$\Omega^{-}_{U}=\Omega^{-}_{U'}$

Figure 3: Scheme of the properties concerning to two observers Ω-related



Figure 4: Scheme of two observers that are not Ω-related

(i) $U \overset{\Omega}{\approx} U'$, *i.e.* U *and* U' *are* Ω*-related.*

(ii) $U \overset{\Omega_U^-}{\approx} U'$, *i.e.* U *and* U' *are* Ω_U^-*-related.*

(iii) $\Omega_U^- = \Omega_{U'}^-$.

(iv) U, U' *and the lightlike direction of* Ω *are in the same* 2*-plane.*

(v) $\Omega \cap U^\perp = \Omega \cap (U')^\perp$.

(vi) $U' \in \left(\Omega \cap U^\perp\right)^\perp$.

(vii) $\Omega_{UU'}^{--} = \Omega_{U'U}^{--} = \Omega$.

Proof.

All the equivalences can be easily proved, except for the property

$$\Lambda_{UU'}^{--} = \Lambda_{U'U}^{--} \longrightarrow U \overset{\Omega}{\approx} U', \tag{9}$$

where Λ is a lightlike 1-distribution (the case of dimension $p = n-1$ can be proved taking into account (7) and (9)).

Let $\{S+U\}$ be the U-basis of Λ and let $\{X_1, \ldots, X_{n-2}, S, U\}$ be an orthonormal basis.

- We are going to build the U'-basis of $\Lambda_{UU'}^{--}$:

  We have that $\{-S+U\}$ is the U-basis of Λ_U^-. If $\{-Y'+U'\}$ is the U'-basis of Λ_U^-, then $\{Y'+U'\}$ is the U'-basis of $\Lambda_{UU'}^{--}$. We can write $U' = \gamma(U+vX)$ (see (5)), and by using algebraic manipulations, we have that

$$\begin{aligned} Y'+U' &= 2v\gamma g(X,X_1)X_1 \\ &+ \ldots + 2v\gamma g(X,X_{n-2})X_{n-2} \\ &+ \left(2v\gamma g(X,S) + \frac{1}{\gamma(1+vg(X,S))}\right)S \\ &+ \left(2\gamma - \frac{1}{\gamma(1+vg(X,S))}\right)U. \end{aligned} \tag{10}$$

- We are going to build the U-basis of $\Lambda_{U'U}^{--}$:

  Let $\{-S'+U'\}$ be the U'-basis of $\Lambda_{U'}^-$ and let $\left\{X_1', \ldots, X_{n-2}', S', U'\right\}$ be an orthonormal basis. If $\{-Y+U\}$ is the U-basis of $\Lambda_{U'}^-$, then $\{Y+U\}$ is the U-basis of $\Lambda_{U'U}^{--}$. Taking into account expression (5) and by using algebraic manipulations, we have that

$$Y+U = -\frac{2v}{2-\alpha}g(X,X_1)X_1$$

$$
\begin{aligned}
& - \ldots - \frac{2v}{2-\alpha} g\left(X, X_{n-2}\right) X_{n-2} \\
& + \left(\frac{\alpha - 2vg\left(X,S\right)}{2-\alpha}\right) S + U, \qquad (11)
\end{aligned}
$$

where $\alpha = \frac{1}{\gamma^2(1-vg(X,S))}$.

By the hypothesis, the vector fields $Y' + U'$ and $Y + U$ are proportional. So, identifying the coefficients of U in (10) and (11) we obtain that

$$Y' + U' = \left(2\gamma - \frac{1}{\gamma\left(1 + vg\left(X,S\right)\right)}\right)\left(Y + U\right). \qquad (12)$$

If we suppose $g\left(X,S\right) \neq \pm 1$ (i.e. $\left|g\left(X,S\right)\right| < 1$), then $g\left(X, X_i\right) \neq 0$ for all $i = 1, \ldots, n-2$. So, identifying the coefficients of X_i in (10) and (11), taking into account (12), we obtain that

$$2v\gamma = \left(2\gamma - \frac{1}{\gamma\left(1 + vg\left(X,S\right)\right)}\right)\left(\frac{-2v}{2-\alpha}\right). \qquad (13)$$

Solving $g\left(X,S\right)$ in (13), with $v \neq 0$

$$g\left(X,S\right) = \pm\sqrt{\frac{v^2+1}{2v^2}},$$

but $v^2 < 1$, so $\frac{v^2+1}{2v^2} > 1$ and then there is not any solution with $\left|g\left(X,S\right)\right| < 1$.

On the other hand, it can be easily proved that if we suppose $g\left(X,S\right) = \pm 1$ (i.e. $X = \pm S$), then $Y' + U'$ and $Y + U$ are always proportional. In this case, taking into account expression (5) we have that U' can be expressed as a linear combination of U and $S + U$. So, $U \overset{\Omega}{\approx} U'$ (by (iv)).

■

5 Equality between associated bases

We can wonder if given a lightlike distribution (of dimension or co-dimension 1) and given two different observers, U and U', it is possible for a U-basis to coincide with a U'-basis of this distribution. First, we are going to study it in the case of dimension 1:

Let $\{S + U\}$ be the U-basis of a lightlike 1-distribution Λ. We are going to build all the different observers U' with U'-basis of Λ equal to the U-basis of Λ. Taking into account (5) we have to find X and v such that

$$g\left(X,S\right) = \frac{1-\sqrt{1-v^2}}{v}. \qquad (14)$$

The function $\frac{1-\sqrt{1-v^2}}{v} :]0,1[\to]0,1[$ is bijective. So, given a spacelike unitary vector field X orthogonal to U such that $0 < g\left(X,S\right) < 1$, there exists a unique v such that (14) is satisfied. Then U' is given by (5). So, we have a different U' for each different X which satisfies all these conditions.

Example 6 *In the Minkowski space-time, expressing the metric g in rectangular coordinates, we consider the lightlike 1-distribution Λ generated by*

$$\left\{\frac{\partial}{\partial z}+\frac{\partial}{\partial t}\right\}. \tag{15}$$

If we consider $U=\frac{\partial}{\partial t}$ as an observer, then (15) is the U-basis of Λ, of the form $\{S+U\}$ where $S=\frac{\partial}{\partial z}$. If we consider, for instance, the spacelike unitary vector field $X=\frac{1}{\sqrt{2}}\left(\frac{\partial}{\partial z}+\frac{\partial}{\partial x}\right)$ orthogonal to U, we have that $g(X,S)=\frac{1}{\sqrt{2}}\in\,]0,1[$ and then, we have that $v=\frac{2\sqrt{2}}{3}$ from (14). So, given the observer

$$U'=\frac{1}{\sqrt{1-v^2}}(U+vX)=3\frac{\partial}{\partial t}+2\left(\frac{\partial}{\partial z}+\frac{\partial}{\partial x}\right),$$

we have that the U'-basis of Λ is the same as the U-basis of Λ.

In the case of co-dimension 1, we need the concept of "observers Ω-related", introduced in Sec. 4. Given Ω a lightlike $(n-1)$-distribution and given U an observer with $\{X_1,\dots,X_{n-2},S+U\}$ a U-basis of Ω, we want to find an observer U' with U'-basis of Ω equal to $\{X_1,\dots,X_{n-2},S+U\}$. If a U'-basis of Ω is given by $\left\{X_1',\dots,X_{n-2}',S'+U'\right\}$, we must impose two conditions:

- $S+U=S'+U'$
- Since we have a certain freedom in the choice of $X_1',\dots,X_{n-2}'$, we have to impose only that $span(X_1,\dots,X_{n-2})=span\left(X_1',\dots,X_{n-2}'\right)$, i.e. $\Omega\cap U^{\perp}=\Omega\cap(U')^{\perp}$.

The first condition is equivalent (according to the case of dimension 1) to the existence of a spacelike unitary vector field X orthogonal to U such that $0<g(X,S)<1$.

The second condition is equivalent (according to Theorem 5 (v)) to $U\overset{\Omega}{\approx}U'$. Then, S,U,U' are in the same $(n-2)$-plane, i.e. U' can be expressed as a linear combination of S and U. Since $U'=\gamma(U+vX)$, we obtain that X can be also expressed as a linear combination of S and U. But X is orthogonal to U and then, the unique possibility is $X=\pm S$ (since they are unitary vector fields). So, $g(X,S)=\pm1$, and this is not possible, according to the first condition.

So, given two different observers, U and U', and a lightlike $(n-1)$-distribution Ω, we have that a U-basis of Ω is always different from any U'-basis of Ω.

6 Conditions of integrability

In this paper, we have studied some properties of lightlike $(n-1)$-distributions in general, which only in some cases are foliations (however, since every vector field is integrable, given any 1-distribution it is automatically a 1-foliation).

Now we are going to study when we can integrate a lightlike $(n-1)$-distribution (see Theorem 8).

It is well known that for any given observer U, its local rest space $U^{\perp}$ is a foliation if and only if U is synchronizable [4]. Taking this into account and given a lightlike $(n-1)$-distribution Ω, we are going to study sufficient conditions for the integrability of the spacelike $(n-2)$-distribution $\Omega \cap U^{\perp}$ in next proposition:

Proposition 7 *Let Ω be a lightlike $(n-1)$-distribution,*

(i) if Ω is a foliation, then $\Omega \cap U^{\perp}$ is also a foliation for any synchronizable observer U.

(ii) if there exists a synchronizable observer U such that $\left(\Omega \cap U^{\perp}\right) \oplus span\,(U)$ is a foliation, then $\Omega \cap U^{\perp}$ is a foliation.

Proof.

(i) given U a synchronizable observer, let $\{X_1, \ldots, X_{n-2}, S+U\}$ be a U-basis of Ω.

Since Ω is a foliation, we have that $[X_i, X_j] \in \Omega$ for all $i, j = 1, \ldots, n-2$.

Moreover, $U^{\perp}$ is also a foliation (because U is synchronizable), so $[X_i, X_j] \in U^{\perp}$ for all $i, j = 1, \ldots, n-2$, and then

$$[X_i, X_j] \in \Omega \cap U^{\perp},$$

for all $i, j = 1, \ldots, n-2$. So, $\Omega \cap U^{\perp}$ is a foliation.

(ii) let $\{X_1, \ldots, X_{n-2}, S+U\}$ be a U-basis of Ω.

Since $\left(\Omega \cap U^{\perp}\right) \oplus span\,(U)$ is a foliation, we have that $[X_i, X_j] \in \left(\Omega \cap U^{\perp}\right) \oplus span\,(U)$ for all $i, j = 1, \ldots, n-2$.

Moreover, $U^{\perp}$ is also a foliation, so $[X_i, X_j] \in U^{\perp}$ for all $i, j = 1, \ldots, n-2$, and then

$$[X_i, X_j] \in \left(\left(\Omega \cap U^{\perp}\right) \oplus span\,(U)\right) \cap U^{\perp} = \Omega \cap U^{\perp},$$

for all $i, j = 1, \ldots, n-2$. So, $\Omega \cap U^{\perp}$ is a foliation.

■

Nevertheless, the integrability of the timelike $(n-1)$-distribution $\left(\Omega \cap U^{\perp}\right) \oplus span\,(U)$ depends exclusively on the integrability of Ω_U^-, as we see in next theorem:

Theorem 8 *Let Ω be a lightlike $(n-1)$-foliation and U a synchronizable observer. Ω_U^- is a foliation if and only if $\left(\Omega \cap U^{\perp}\right) \oplus span\,(U)$, is a foliation.*

Proof.

Let $\{X_1, \ldots, X_{n-2}, S+U\}$ be a U-basis of Ω.

By Proposition 7 (i) we have that $\Omega \cap U^{\perp}$ is a foliation.

$\longrightarrow$ Moreover, taking into account that Ω and Ω_U^- are foliations, it can be proved that $[X_i, U] \in \mathit{span}\,(X_1, \ldots, X_{n-2}, U) = \left(\Omega \cap U^\perp\right) \oplus \mathit{span}\,(U)$ for all $i = 1, \ldots, n-2$, and so $\left(\Omega \cap U^\perp\right) \oplus \mathit{span}\,(U)$ is a foliation.

$\longleftarrow$ Taking into account that Ω and $\left(\Omega \cap U^\perp\right) \oplus \mathit{span}\,(U)$ are foliations, it can be proved that $[X_i, -S + U] \in \Omega_U^-$ for all $i = 1, \ldots, n-2$, and so Ω_U^- is a foliation.

■

Theorem 8 assures that the observation of the foliation Ω_U^- is guaranteed only when $\left(\Omega \cap U^\perp\right) \oplus \mathit{span}\,(U)$ is a foliation. Note that Theorem 8 does not state that if Ω is a lightlike $(n-1)$-foliation and U is a synchronizable observer, then Ω_U^- is a foliation. Actually we show with an example that this statement is false:

Example 9 *In the Minkowski space-time, expressing the metric g in rectangular coordinates, we consider the following vector fields:*

$$\begin{aligned} U &= \frac{\partial}{\partial t} + a\frac{\partial}{\partial x} + b\frac{\partial}{\partial y} + c\frac{\partial}{\partial z} = (1, a, b, c) \\ X_1 &= a\frac{\partial}{\partial t} + \frac{\partial}{\partial x} = (a, 1, 0, 0) \\ X_2 &= b\frac{\partial}{\partial t} + \frac{\partial}{\partial y} = (b, 0, 1, 0) \\ S &= \frac{c}{d}\left(\frac{\partial}{\partial t} + a\frac{\partial}{\partial x} + b\frac{\partial}{\partial y} + \frac{\alpha^2}{c}\frac{\partial}{\partial z}\right) = \frac{c}{d}\left(1, a, b, \frac{\alpha^2}{c}\right) \end{aligned}$$

where a, b, c, d are smooth functions such that $c^2 < 1 - \left(a^2 + b^2\right) = d^2$, U is an observer and $\{X_1, X_2, S\}$ are spacelike vector fields defining an orthogonal basis of $U^\perp$. We consider Ω the 3-distribution generated by $\{X_1, X_2, S + U\}$ where $S + U = \left(1 + \frac{c}{d}\right)(1, a, b, d)$. Stating $b = 0$, defining d by the equation $\frac{x}{d\sqrt{1-d^2}} - \frac{z}{(f+d^3)^{2/3}} = 2$, and taking

$$f = \left(\frac{1}{4}\right)^{3/2} - \left(\frac{1}{2}\right)^{3/2}, \quad c = \left(f + d^3\right)^{1/3}, \quad a^2 = 1 - d^2,$$

we obtain that in a neighborhood of $W := \{(t, x, y, z) : x = 1, z = 0\} \subset M$, U is synchronizable, Ω is a foliation and Ω_U^- is not a foliation.

Moreover, Example 9 ensures that conditions (i) and (ii) of Proposition 7 are not necessary conditions for the integrability of $\Omega \cap U^\perp$:

(i) Ω_U^- is not a foliation, but $\Omega_U^- \cap U^\perp$ is a foliation.

(ii) U is synchronizable and $\left(\Omega \cap U^\perp\right) \oplus \mathit{span}\,(U)$ is not a foliation, but $\Omega \cap U^\perp$ is a foliation.

7 Some physical interpretations and applications

According to Sec. 1, a congruence of lightlike rays (rays of light for example) are, in fact, the leaves of a lightlike 1-foliation Λ. On the other hand, a congruence of lightlike moving wave fronts are the leaves of a lightlike $(n-1)$-foliation Ω. Precisely, given an observer U, the leaves of $\Omega \cap U^{\perp}$ can be interpreted as the spacelike wave fronts ($(n-2)$-dimensional without movement) for this observer [3]. But neither all lightlike 1-foliations can be interpreted as a congruence of lightlike rays nor all lightlike $(n-1)$-foliations can be interpreted as a congruence of lightlike moving wave fronts. For example, Λ and Ω should be totally geodesic foliations. Moreover, in the case of dimension $(n-1)$, condition (i) of Proposition 7 should be a sufficient and necessary condition: Ω is integrable if and only if $\Omega \cap U^{\perp}$ is integrable (given a synchronizable observer U). This fact is reasonable, since the observation of the wave fronts (as leaves of $\Omega \cap U^{\perp}$) should imply the existence of the wave fronts (as leaves of Ω) and viceversa. In this Section, we are going to work in a 4-dimensional space-time (i.e. $n=4$) with foliations that can be interpreted in this way (as congruences of lightlike rays or lightlike moving wave fronts) to make physical applications of some results given in this paper.

7.1 Light aberration

Let Ω be a lightlike foliation of dimension 1 or 3, and let U, U' be two observers. If N represents the lightlike direction of Ω, let S, S' represent the relative direction of N for the observers U, U', respectively (i.e. $S+U$ and $S'+U'$ are proportional to N). These relative directions are the spacelike directions of propagation of the lightlike rays (dimension 1) or the lightlike moving wave fronts (dimension 3) for these observers, respectively. So, we have that

- if $U \overset{\Omega}{\approx} U'$ (i.e. they are Ω-related), then S and S' are proportional. In this case, they represent the same relative direction.
- if U and U' are not Ω-related, then S, S' represent different relative directions. As we said in Sec. 3, this fact is the light aberration effect.

  Taking into account (6) we can obtain, for example, the usual expression for light aberration (see [6]) in a more general way:

  $$\cos\theta = \frac{\cos\theta' - v}{1 - v\cos\theta'},$$

  where θ is the angle between $-S$ and X, (i.e. $\cos\theta = g(X, -S)$) and θ' is the angle between $-S'$ and the projection of X to $U'^{\perp}$.

7.2 Stationary waves

It is known that a stationary wave is formed in fact by two identical waves with opposite relative directions of propagation. So, a 3-dimensional lightlike stationary wave for an observer U can be represented as a pair of lightlike 3-foliations $\left\{\Omega, \Omega_U^-\right\}$ (actually they represent the 3-dimensional wave fronts), since the foliations Ω and Ω_U^- have opposite relative directions of propagation for the observer U. So, a 3-dimensional lightlike stationary wave for an observer U is represented by an equivalence class of the relation "be equal up to orientations for the observer U", given in Definition 3 and in expression (8). In this case, if U and U' are two observers Ω-related, $\left\{\Omega, \Omega_U^-\right\}$ represents a stationary wave for both observers U and U'.

Finally, as we are only interested in lightlike 3-foliations Ω with physical interpretation (representing congruences of lightlike moving wave fronts), condition (i) of Proposition 7 should be a sufficient and necessary condition (this fact is discussed at the beginning of this Section). So, given U a synchronizable observer, Ω_U^- should be also a foliation, since $\Omega \cap U^\perp = \Omega_U^- \cap U^\perp$ is a foliation. If we take this into account, Example 9 does not have any physical interpretation, since Ω cannot represent any congruence of lightlike moving wave fronts. In fact, Ω is not a totally geodesic foliation.

References

[1] V. J. Bolós. Estudio de Foliaciones en Relatividad (Publicacions de la Universitat de València: Valencia 2003).

[2] F. De Felice, C. J. S. Clarke. Relativity on Curved Manifolds (Cambridge University Press: Cambridge 1990).

[3] V. Liern, J. Olivert. A symplectic interpretation of some wave fronts in general relativity. *J. Math. Phys.* **36 (2)** (1995), 837.

[4] R. K. Sachs, H. Wu. Relativity for Mathematicians (Springer Verlag: Berlin, Heidleberg, New York 1977).

[5] J. M. Souriau. Structure of Dynamical Systems. A Symplectic View of Physics (Birkhäuser: Boston 1997).

[6] J. L. Synge. Relativity: The Special Theory (North-Holland Publishing: Amsterdam 1965).